



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$x = \frac{\begin{vmatrix} .24561, & .35381, & .36518 \\ .62433, & .90274, & .91143 \\ .17145, & .23642, & .24375 \end{vmatrix}}{\begin{vmatrix} .34622, & .35381, & .36518 \\ .89318, & .90274, & .91143 \\ .22431, & .23642, & .24375 \end{vmatrix}}.$$

If in the numerator determinant two thirds of the third column is subtracted from the first column, and if the second column is subtracted from the third, the value of the determinant is not altered. In the denominator let the middle column be subtracted from each of the others, and then the third column added to the first. The resulting expression for x is

$$x = \frac{\begin{vmatrix} .00216, & .35381, & .01137 \\ .01671, & .90274, & .00869 \\ .00895, & .23642, & .00733 \end{vmatrix}}{\begin{vmatrix} +.00378, & .35381, & .01137 \\ -.00087, & .90274, & .00869 \\ -.00478, & .23642, & .00733 \end{vmatrix}}.$$

The computation can now be made with four-place tables containing all the numbers required on two pages. In computing x it is found in two or three minutes that to three figures its value is -1.03 , which is as far as it is determinate.

THE UNIVERSITY OF CHICAGO,
May 19, 1913.

THE ACCURACY OF INTERPOLATION IN A FIVE-PLACE TABLE OF LOGARITHMS OF SINES.

By A. M. KENYON and G. JAMES, Purdue University.

In a table which gives the logarithms of sines for each minute of angle, the logarithms of sines of angles to tenths of minutes are obtained by interpolation.

The name *Ordinary Interpolation* will be used to denote the process based on the assumption:

(1) As the angle increases from m' to $(m+1)'$ the increment of the logarithm of its sine varies as the increment of the angle; that is,

$$\log \sin (m + n/10)' = \log \sin m' + (n/10)(\log \sin(m+1)' - \log \sin m') \\ (n = 1, 2, 3, \dots 9).$$

The error introduced by ordinary interpolation arises from two sources:

(a) The *essential* error, due to the assumption (1).

(b) The *tabular* error, due to the fact that the tabulated values of $\log \sin m'$ and of $\log \sin(m+1)'$ are in error.

The essential error increases as the angle decreases and makes ordinary interpolation unreliable for small angles. The tabular error is irregular but nowhere in the table does it exceed, in absolute value, one unit in the fifth place.

To make this clear we need to note that whenever an irrational number is approximated by a rational number correct to n decimal places, an error is introduced which in absolute value is less than $5/10^{n+1}$. Thus if we write $\sin 45^\circ = .70711$, the result is too large by an error between .000003 and .000004.

Let $\log \sin m' = a$, $\log \sin(m + 1)' = b$ and let $a - e_1$, $b - e_2$, be the five place values of these same logarithms. Then by ordinary interpolation (assumed to be a correct process since we seek here only the tabular error), the interpolated value of $\log \sin(m + n/10)'$ would be

$$c = a + (n/10)(b - a) = c_1 - e_3$$

where c_1 represents the five place value of c .

But by ordinary interpolation from a five place table, the result will be

$$\begin{aligned} c_2 &= a - e_1 + (n/10)(b - e_2 - a + e_1) \\ &= a + (n/10)(b - a) - [e_1 + (n/10)(e_2 - e_1)] \\ &= c_1 - [e_3 + e_1 + (n/10)(e_2 - e_1)]. \end{aligned}$$

Whence the five place tabular error is

$$c_1 - c_2 = (\bar{n}/10)e_1 + (n/10)e_2 + e_3, \text{ where } n + \bar{n} = 10,$$

and obviously the absolute value of this error is less than .00001.

Actual trial of a very large number of cases (over 1,200) shows that ordinary interpolation below 2° gives the result correct to five places in less than 30 per cent. of all cases; that it gives an error of one in the fifth place in about 50 per cent., and an error of two or more in the fifth place in about 20 per cent. of cases. There is no case above $1^\circ 30'$ where the total error is greater than one unit in the fifth place. Above 2° it gives the result correct to five places in from 70 per cent. to 83 per cent. of all cases and in no case tried is the error greater than one unit in the fifth place.

The name *Logarithmic Interpolation* will be used to denote the process based on the following assumption:

(2) For sufficiently small angles, the sine varies as the angle; which gives, if m and n denote measures of angles in the same unit,

$$\log \sin(m + n/10) = \log \sin m + \log(m + n/10) - \log m.$$

Here the essential error decreases with the angle; the tabular error as before is irregular but nowhere in the table exceeds, in absolute value, one unit in the fifth place.

Actual trial of all cases (to tenths of minutes) shows that in no case below $5^\circ 45'$ does logarithmic interpolation introduce a total error greater than one in the

fifth place.¹ Below 5° it gives the result correct to five places in about 60 per cent. of all cases and shows an error of one in the fifth place in about 40 per cent. of cases.

Below 2° , logarithmic interpolation is certainly preferable to ordinary interpolation for three reasons: (1) it gives the correct result in a larger number of cases; (2) it gives an error in the fifth place in a smaller number of cases; (3) it never gives an error greater than one in the fifth place.

Above 2° , ordinary interpolation is preferable for the same three reasons. All cases between $1^\circ 30'$ and $2^\circ 15'$ were examined.

An examination of 200 cases in the neighborhood of 84° shows that ordinary interpolation will give the result correct to five places in about 83 per cent., and will show an error of one in the fifth place in 17 per cent. of cases.

Below is a table of results of the statistical investigation.

Interval of Angle.	Number of Cases Examined.	Per Cent. of Cases Giving Result Correct to 5 Places.		Per Cent. of Cases Showing an Error of One in the 5th Place.		Per Cent. of Cases Showing an Error of Two or More in the Fifth Place.	
		Ordinary.	Logarithmic.	Ordinary.	Logarithmic.	Ordinary.	Logarithmic. ²
$1'-1^\circ$	45	Fails	60	Large	40		0
$1^\circ -1^\circ 30'$	97	33	57	57		10	0
$1^\circ 30'-1^\circ 45'$	All	55	63	45	37	0	0
$1^\circ 45'-2^\circ$	All	63	69	37	31	0	0
$2^\circ -2^\circ 15'$	All	71	62	29	38	0	0
$2^\circ 15'-3^\circ$	55	78	60	22	40	0	0
$3^\circ -4^\circ$	55	80		20		0	0
$4^\circ -5^\circ$						0	0
$5^\circ -5^\circ 45'$	All		66		34	0	0
$5^\circ 45'-6^\circ$	All		64		36		*

A THEOREM ABOUT ISOGONAL CONJUGATES.

By DAVID F. BARROW, Harvard University.

As an introduction let us recall two well-known theorems of elementary plane geometry:

THEOREM I. *Given a triangle $A_1A_2A_3$ (Fig. 1) and any point P not a vertex, join P to the three vertices and reflect each of the three lines thus drawn in the bisector of the angle at the corresponding vertex. The three reflected lines will meet in a point P' , called the isogonal conjugate of P .*

THEOREM II. *Given a triangle and a point marked at random on each side. If three circles be drawn, one through each vertex and the two adjacent marked points, these three circles meet in a common point.*

¹ The first error of two in the fifth place enters at $5^\circ 53'.7$

² All cases examined for this column.

* At $5^\circ 53'.7$ the first error of 2 in the fifth place enters.